

B2x  
18. The method according to claim 13, wherein the daily dose of formoterol, including maintenance therapy, lies in the range of from 1  $\mu\text{g}$  to 100  $\mu\text{g}$ , calculated as formoterol fumarate dihydrate.

---

B3  
20. The method according to claim 13, wherein a unit dose of budesonide lies in the range of from 20  $\mu\text{g}$  to 1600  $\mu\text{g}$ .

X  
21. The method according to claim 13, wherein the daily dose of budesonide, including maintenance therapy, lies in the range of from 20  $\mu\text{g}$  to 4800  $\mu\text{g}$ --

---

Please add claims 25-34.

---

B4  
-- 25. The method according to claim 14, wherein the molar ratio of (a) to (b) calculated as formoterol to budesonide is from 1:1 to 1:70.

26. The method according to claim 17, wherein a unit dose of formoterol lies in the range of from 3  $\mu\text{g}$  to 12  $\mu\text{g}$ , calculated as formoterol fumarate dihydrate.

27. The method according to claim 18, wherein the daily dose of formoterol, including maintenance therapy, lies in the range of from 2  $\mu\text{g}$  to 60  $\mu\text{g}$ , calculated as formoterol fumarate dihydrate.  
6 - 48  $\mu\text{g}$

X  
28. The method according to claim 20, wherein a unit dose of budesonide lies in the range of from 50  $\mu\text{g}$  to 400  $\mu\text{g}$ .  
50 - 4800  $\mu\text{g}$   
100 - 16000  $\mu\text{g}$

29. The method according to claim 21, wherein the daily dose of budesonide, including maintenance therapy, lies in the range of from 30  $\mu\text{g}$  to 3200  $\mu\text{g}$ .

30. The method according to claim 13 further comprising instructing the patient to inhale the composition as a rescue medication.

31. The method according to claim 13 further comprising instructing the patient to take a second composition, comprising a glucocorticosteroid, on a regular basis as a maintenance treatment.

32. The method according to claim 13 further comprising instructing the patient to use the composition as a complement to maintenance treatment of the patient's asthma.

33. The method according to claim 13 further comprising instructing the patient to inhale an effective amount of the composition as a preventive measure prior to encountering an asthma triggering event.

34. The method of claim 33 wherein the asthma triggering event is selected from the group consisting of exposure to cold air, exercise, and exposure to a smoky environment.--

---